

Shearing coordinates and convexity of length functions on Teichmüller space

M. Bestvina, K. Bromberg, K. Fujiwara, J. Souto*

February 5, 2009

Abstract

We prove that there are Fenchel-Nielsen coordinates for the Teichmüller space of a finite area hyperbolic surface with respect to which the length functions are convex.

1 Introduction

Let X be a complete hyperbolic surface with finite area and $\mathcal{T}(X)$ its Teichmüller space. Recall that points in $\mathcal{T}$ are equivalence classes of marked hyperbolic surfaces, i.e. equivalence classes of pairs (Y, f) where Y is a hyperbolic surface and $f : X \rightarrow Y$ is a quasi-conformal homeomorphism.

The goal of this paper is to describe certain coordinates of $\mathcal{T}(X)$ with respect to which the length functions are convex. Given a homotopically essential, non-peripheral curve $\gamma \subset X$ and a point $(Y, f) \in \mathcal{T}(S)$, then the curve $f(\gamma)$ is freely homotopic to a unique geodesic in the hyperbolic surface Y . Denoting by $\ell_\gamma(Y, f)$ the length of this geodesic we obtain a well-defined function

$$\ell_\gamma : \mathcal{T}(X) \rightarrow \mathbb{R}_+$$

This is the *length function* associated to the curve γ .

Of the many well-known ways to parametrize Teichmüller space perhaps the most classical is attributed to Fenchel and Nielsen. Fix a pants decomposition $\mathcal{P}$ of X , i.e. a multicurve such that $X \setminus \mathcal{P}$ is homeomorphic to the disjoint union of thrice punctured

*M.B. has been partially supported by NSF grant DMS-0502441, K.B. has been partially supported by NSF grants DMS-0554569 and DMS-0504877, K. F. has been partially supported by Grant-in-Aid for Scientific Research (No. 19340013) and J. S. has been partially supported by NSF grant DMS-0706878 and the Alfred P. Sloan Foundation.

spheres. A simple computation shows that $\mathcal{P}$ has $|\mathcal{P}| = 3g + n - 3$ components where g is the genus and n the number of cusps of X . The Fenchel-Nielsen coordinates

$$\Phi : \mathcal{T}(X) \xrightarrow{\sim} \mathbb{R}_+^{3g+n-3} \times \mathbb{R}^{3g+n-3} \quad (1.1)$$

associate to each point in $\mathcal{T}(X)$ the length and the twist for each component of the pants decomposition $\mathcal{P}$.

In the above discussion the meaning of "length" is clear. To define the "twist parameter" it has to be decided what a 0-twist represents. There is no canonical way to determine this so we refer to any of the possible length-twist coordinates as *Fenchel-Nielsen coordinates* associated to the pants decomposition $\mathcal{P}$. We prove:

Theorem 1.1 *Let X be a complete, finite area, hyperbolic surface of genus g and with n cusps, and fix a pants decomposition $\mathcal{P}$ of X . There are Fenchel-Nielsen coordinates $\Phi : \mathcal{T}(X) \xrightarrow{\sim} \mathbb{R}_+^{3g+n-3} \times \mathbb{R}^{3g+n-3}$ associated to $\mathcal{P}$ such that for any essential curve γ in X the function*

$$l_\gamma \circ \Phi^{-1} : \mathbb{R}_+^{3g+n-3} \times \mathbb{R}^{3g+n-3} \rightarrow \mathbb{R}_+$$

is convex. If moreover the curve γ intersects all the components of $\mathcal{P}$ then $l_\gamma \circ \Phi$ is strictly convex.

Other convexity results of the length functions l_γ are due to Kerckhoff [7] and Wolpert [8]. They proved respectively that the length functions are convex along earthquake paths and Weil-Petersson geodesics. Both authors derived from their results proofs of the so-called Nielsen realization problem; so do we.

Theorem 1.2 (Kerckhoff) *The action of every finite subgroup of the mapping class group on $\mathcal{T}(X)$ has a fixed point.*

Tromba [10] gave a different proof of Theorem 1.2 using the convexity of the energy functional along Weil-Petersson geodesics. Proofs of this theorem in a completely different spirit are due to Gabai [5] and Casson-Jungreis [4].

In order to prove Theorem 1.1 we follow a slightly indirect path. We will associate a continuous map $s_\lambda : \mathcal{T}(X) \rightarrow \mathbb{R}^{|\lambda|}$ to every maximal, finite leaved lamination λ of X . The image $T_\lambda = s_\lambda(\mathcal{T}(X))$ of s_λ is an open convex subset of a linear subspace of the correct dimension $6g + 2n - 6$. We refer to

$$s_\lambda : \mathcal{T}(X) \xrightarrow{\sim} T_\lambda$$

as *shearing coordinates* associated to the lamination λ . The shearing coordinates are closely related to those studied by Bonahon [1].

Given a pants decomposition $\mathcal{P}$ of X , we choose a maximal lamination λ containing $\mathcal{P}$ and describe Fenchel-Nielsen coordinates associated to $\mathcal{P}$ in such a way that the map

$$\Phi \circ s_\lambda^{-1} : T_\lambda \xrightarrow{\sim} \mathbb{R}_+^{3g+n-3} \times \mathbb{R}^{3g+n-3}$$

is linear. In particular, Theorem 1.1 follows immediately from the following more general result:

Theorem 1.3 *Let X be a complete, finite area, hyperbolic surface with genus g and with n cusps. Let λ be a maximal lamination in X with finitely many leaves and let $s_\lambda : \mathcal{T}(X) \xrightarrow{\sim} T_\lambda$ be the shearing coordinates associated to λ . For any essential curve γ in X the function*

$$l_\gamma \circ s_\lambda^{-1} : T_\lambda \rightarrow \mathbb{R}_+$$

is convex. If moreover the curve γ intersects all the leaves of λ then $l_\gamma \circ s_\lambda^{-1}$ is strictly convex.

The paper is organized as follows. After a few preliminaries in section 2, we introduce in section 3 the shearing coordinates and reduce the proof of Theorem 1.3 to Proposition 3.6, our main technical result. In section 4 we relate Fenchel-Nielsen coordinates to shearing coordinates and prove Theorem 1.1. In section 5 we study the length function on the Teichmüller space of the annulus and finally in section 6 we prove Proposition 3.6. The proofs are, once one is used to the notation, elementary.

2 Teichmüller space and other important objects

Let X be a complete, orientable hyperbolic surface with possibly infinite area. A *marked hyperbolic surface* is a pair (Y, f) where Y is a hyperbolic surface and $f : X \rightarrow Y$ is a quasi-conformal homeomorphism. Two marked surfaces (Y_0, f_0) and (Y_1, f_1) are equivalent if there is an isometry $\phi : Y_0 \rightarrow Y_1$ such that $\phi \circ f_0$ and f_1 are quasi-conformally isotopic. The Teichmüller space $\mathcal{T}(X)$ is the set of equivalence classes of marked hyperbolic surfaces. We give $\mathcal{T}(X)$ a metric (and topology) as follows. The distance, $d_T((Y_0, f_0), (Y_1, f_1))$, between two pairs is the infimum of the logarithm of the quasi-conformal constant of all maps $\phi : Y_0 \rightarrow Y_1$ with $\phi \circ f_0$ quasi-conformally isotopic to f_1 .

Remark. In this paper we will be mostly interested in Teichmüller spaces of finite area surfaces although the ambitious reader could easily see that our results can be extended to a more general setting. In particular, as we will see in section 5 the Teichmüller space of a hyperbolic annulus plays a key role in our work.

Continuing with the same notation as above, we can identify the universal covers of X and Y with the hyperbolic plane $\mathbb{H}^2$; this identification is unique up to composition

with an isometry of $\mathbb{H}^2$. Let $\tilde{f} : \mathbb{H}^2 \rightarrow \mathbb{H}^2$ be the lift of the quasi-conformal homeomorphism $f : Y \rightarrow X$. It is well-known that $\tilde{f}$ extends continuously to a homeomorphism of $\partial\tilde{f} : \partial\mathbb{H}^2 \rightarrow \partial\mathbb{H}^2$; here $\partial\mathbb{H}^2$ is the boundary at infinity of $\mathbb{H}^2$. Moreover, lifts of quasi-conformally isotopic maps have extensions which differ by composition with (the boundary extensions of) isometries of $\mathbb{H}^2$.

A *lamination* on X is a closed, but perhaps not compact, subset of X which is foliated by geodesics. If the surface X has finite area, we will be only be interested in laminations λ with only finitely many leaves. Recall that the only non-isolated leaves of such a lamination are simple closed geodesics in X .

Let $f : X \rightarrow Y$ be a quasi-conformal homeomorphism between hyperbolic surfaces and let $\tilde{f} : \mathbb{H}^2 \rightarrow \mathbb{H}^2$ and $\partial\tilde{f} : \partial\mathbb{H}^2 \rightarrow \partial\mathbb{H}^2$ be as above. The maps f and $\tilde{f}$ will not take geodesics to geodesics. In order to by-pass this problem, we associate to any geodesic $\gamma \subset \mathbb{H}^2$ the unique geodesic $\bar{f}(\gamma) \subset \mathbb{H}^2$ which has the same endpoints on $\partial\mathbb{H}^2$ as the arc $\tilde{f}(\gamma)$. If γ is a closed geodesic on X then its pre-image $\tilde{\gamma}$ in $\mathbb{H}^2$ will be equivariant and therefore $\bar{f}(\tilde{\gamma})$ will also be equivariant and descend to a geodesic $\bar{f}(\gamma)$ on Y . If γ is also simple then $\bar{f}(\gamma)$ is simple as well. For a lamination we apply $\bar{f}$ to each geodesic in the lamination. Similarly, for an ideal triangle Δ we let $\bar{f}(\Delta)$ be the ideal triangle whose boundary is the the $\bar{f}$ -image of the geodesics bounding Δ .

We refer to [6] and [3] for facts and definitions in this section.

3 Shearing coordinates

The goal of this section is to define *shearing coordinates* for the Teichmüller space $\mathcal{T}(X)$ of a finite area hyperbolic surface.

3.1 Ideal triangulations

Before setting up our coordinates we need some definitions and notation. An *ideal triangle* on a hyperbolic surface X is the image of an injective, local isometry from an ideal triangle in $\mathbb{H}^2$ to the surface. Recall that any two ideal triangles in $\mathbb{H}^2$ are isometric and that ideal triangles have the same isometry group as euclidean equilateral triangles. An *ideal triangulation* of X is a lamination with finitely many leaves whose complementary components are ideal triangles. An ideal triangle has a unique inscribed disk that is tangent to all three sides of the triangle. The *midpoints* of the sides are three tangency points; compare with figure 1 (a).

Let γ_0 and γ_1 be geodesics in $\mathbb{H}^2$ that are asymptotic to a point $p_\infty \in \partial\mathbb{H}^2$. For each $p_0 \in \gamma_0$ there is a unique $p_1 \in \gamma_1$ such that the horocycle based at p_∞ through p_0 intersects γ_1 at p_1 . We define a map $h_{\gamma_0, \gamma_1} : \gamma_0 \rightarrow \gamma_1$ by $h_{\gamma_0, \gamma_1}(p_0) = p_1$. We allow the possibility that $\gamma_0 = \gamma_1$ in which case h_{γ_0, γ_1} is the identity map; compare with figure 1 (b).




Figure 1: (a) and (b)

Let Δ^a and Δ^b be ideal triangles in $\mathbb{H}^2$ with disjoint interiors and let γ be a geodesic separating the two triangles. We also assume that both Δ^a and Δ^b are asymptotic to γ ; that is, there are sides γ^a and γ^b of Δ^a and Δ^b that are asymptotic to γ . Let m^a and m^b be the midpoints of γ^a and γ^b .

We define $s(\Delta^a, \Delta^b, \gamma)$ to be the signed distance between $h_{\gamma^a, \gamma}(m^a)$ and $h_{\gamma^b, \gamma}(m^b)$ where the sign is determined by orienting γ such that Δ^a is on the left of γ ; compare with figure 2. If Δ^a and Δ^b have a common boundary edge there is only one choice for γ so we will sometimes write $s(\Delta^a, \Delta^b) = s(\Delta^a, \Delta^b, \gamma)$.




Figure 2: The two dots are $h_{\gamma^a, \gamma}(m^a)$ and $h_{\gamma^b, \gamma}(m^b)$. In this picture $s(\Delta^a, \Delta^b, \gamma)$ is equal to the length of the bold printed segment.

The following lemma is a collection of simple facts on $s(\cdot, \cdot, \cdot)$ whose proof we leave to the interested reader.

Lemma 3.1 *Let Δ^a , Δ^b and γ be as above.*

- $s(\Delta^b, \Delta^a, \gamma) = s(\Delta^a, \Delta^b, \gamma)$.

- If ϕ is an isometry of $\mathbb{H}^2$ then $s(\Delta^a, \Delta^b, \gamma) = s(\phi(\Delta^a), \phi(\Delta^b), \phi(\gamma))$.
- $s(\Delta^a, \Delta^b, \gamma') = s(\Delta^a, \Delta^b, \gamma)$ for any another geodesic γ' separating Δ^a and Δ^b and asymptotic to γ .

If moreover $\Delta^0, \Delta^1, \dots, \Delta^k$ is a chain of pairwise asymptotic ideal triangles with disjoint interior and γ_i is for $i = 1, \dots, k$ a geodesic separating $\Delta^0, \dots, \Delta^{i-1}$ from $\Delta^i, \dots, \Delta^k$ then we have

$$s(\Delta^0, \Delta^k, \gamma) = s(\Delta^0, \Delta^1, \gamma_1) + \dots + s(\Delta^{k-1}, \Delta^k, \gamma_k)$$

for any geodesic γ separating Δ^0 and Δ^k .

3.1

Remark. Working in say the upper half-plane model, assume that two ideal triangles $\Delta_a = [\theta_1, \theta_2, \theta_3]$ and $\Delta_b = [\theta_1, \theta_3, \theta_4]$ have a common boundary edge $[\theta_1, \theta_3]$. Let

$$\kappa = [\theta_1, \theta_2, \theta_3, \theta_4] = \frac{(\theta_1 - \theta_3)(\theta_2 - \theta_4)}{(\theta_1 - \theta_4)(\theta_2 - \theta_3)}$$

be the cross-ratio of the four vertices $\theta_1, \theta_2, \theta_3, \theta_4$. We have then the formula

$$s(\Delta^a, \Delta^b) = \log(\kappa - 1)$$

Decomposing the general picture into adjacent ideal triangles and using the last claim of Lemma 3.1, it is easy to also express $s(\Delta^a, \Delta^b, \gamma)$ as sums of logarithms of algebraic expressions in cross ratios when Δ^a and Δ^b don't have a common boundary edge; compare with figure 3.

3.2 The coordinates

Let λ be from now on an ideal triangulation of the complete finite area hyperbolic surface X . We now define a coordinate map $s_\lambda : \mathcal{T}(X) \longrightarrow \mathbb{R}^{|\lambda|}$ by defining a coordinate function s_γ for each leaf γ of λ . We need to make some choices in the definition of s_λ but the map will be unique up to post-composition with a linear map.

Assume γ is an isolated leaf of λ and let $\tilde{\gamma}$ be a component of the pre-image of γ in the universal cover. Let $\tilde{\lambda}$ be the pre-image of λ in the universal cover. Let Δ_γ^a and Δ_γ^b be the two ideal triangle in the complement of $\tilde{\lambda}$ whose boundary contains $\tilde{\gamma}$. Then we define $s_\gamma(Y, f) = s(\tilde{f}(\Delta_\gamma^a), \tilde{f}(\Delta_\gamma^b), \tilde{f}(\tilde{\gamma}))$. Here $\tilde{f}$ is as in the end of section 2. It is clear that $s_\gamma(Y, f)$ is independent of the choice of the lift $\tilde{\gamma}$.

Now assume that γ is a closed curve in λ . To define s_γ we need to make some arbitrary choices. Again let $\tilde{\gamma}$ be a component of the pre-image of γ in the universal cover. In this case there will not be ideal triangles whose boundary contains $\tilde{\gamma}$. Instead




Figure 3: The dotted lines determine the auxiliary ideal triangles needed to express $s(\Delta^a, \Delta^b, \gamma)$ in terms of cross ratios.

we choose ideal triangles Δ_γ^a and Δ_γ^b such that $\tilde{\gamma}$ separates the triangles and they are both asymptotic to $\tilde{\gamma}$. We then define $s_\gamma(Y, f) = s(\bar{f}(\Delta_\gamma^a), \bar{f}(\Delta_\gamma^b), \bar{f}(\tilde{\gamma}))$.

Remark. Observe that by the last claim of Lemma 3.1, different choices of Δ^a and Δ^b , say $\hat{\Delta}^a$ and $\hat{\Delta}^b$, yield functions $\hat{s}_\gamma(Y, f) = s(\bar{f}(\hat{\Delta}_\gamma^a), \bar{f}(\hat{\Delta}_\gamma^b), \bar{f}(\tilde{\gamma}))$ which differ from $s_\gamma(Y, f)$ by a linear combination of the functions $s_\eta(Y, f)$ corresponding to the isolated leaves η separating Δ^a from $\hat{\Delta}^a$ and Δ^b from $\hat{\Delta}^b$.

We now define our coordinate map by

$$s_\lambda : \mathcal{T}(X) \longrightarrow \mathbb{R}^{|\lambda|} \quad (3.2)$$

by $s_\lambda(Y, f) = (s_{\gamma_1}(Y, f), \dots, s_{\gamma_n}(Y, f))$ where $\gamma_1, \dots, \gamma_n$ are the components of λ . By the remark above, different choices amount to postcomposing s_λ with an invertible linear transformation of $\mathbb{R}^{|\lambda|}$.

To see that s_λ is continuous we fix an identification of $\tilde{X} = \mathbb{H}^2$ with the upper half-plane. Given a quadruple $(\theta_1, \theta_2, \theta_3, \theta_4)$ with $\theta_i \in \partial\mathbb{H}^2 = \mathbb{R} \cup \{\infty\}$ we then define a function

$$\mathcal{T}(X) \rightarrow \mathbb{R}, \quad (Y, f) \mapsto [\partial\tilde{f}(\theta_1), \partial\tilde{f}(\theta_2), \partial\tilde{f}(\theta_3), \partial\tilde{f}(\theta_4)].$$

Since the cross-ratio is invariant under isometries of $\mathbb{H}^2$ this function is well-defined. It directly follows from the geometric definition of a quasi-conformal map that this function is continuous. Since $s_\lambda(Y, f)$ can be expressed in terms of cross-ratios it follows that s_λ is continuous.

3.3 Image of s_λ

Our next goal is to determine the image of s_λ . Before doing so we need still some more notation.

Assume that γ is a non-isolated, i.e. closed, leaf of λ . Our choice of Δ_γ^a and Δ_γ^b determines an a -side and a b -side of γ . In particular if C_γ is a collar neighborhood of γ then $C_\gamma \setminus \gamma$ has two components which we call the *sides* of γ . If $\tilde{C}_\gamma$ is the component of the pre-image of C_γ that contains $\tilde{\gamma}$ then Δ_γ^a will intersect one of the components of $\tilde{C}_\gamma \setminus \tilde{\gamma}$. The image of this component in X will be one of the components of $C_\gamma \setminus \gamma$. This is the a -side. Then Δ_γ^b will intersect the other component of $\tilde{C}_\gamma \setminus \tilde{\gamma}$ and this component will map to the b -side of γ .

We need to assign a sign to each side of γ that will be determined by the direction Δ_γ^a and Δ_γ^b spiral around γ . As above, orient $\tilde{\gamma}$ so that Δ_γ^a is on the left. Then $\sigma_\gamma^a = -1$ if Δ_γ^a is asymptotic to the forward end of $\tilde{\gamma}$ and $\sigma_\gamma^a = +1$ if Δ_γ^a is asymptotic to the negative end. We make a similar definition for σ_γ^b . Note that since $\tilde{\gamma}$ separates Δ_γ^a from Δ_γ^b we need to change the orientation of $\tilde{\gamma}$ when we define σ_γ^b .




Figure 4: In this picture $\sigma_\gamma^a = -1$ and $\sigma_\gamma^b = 1$.

Still assuming that γ is a non-isolated leaf of λ , we will now express its length in terms of our coordinates. The intersection of each leaf β of λ with the a -side of γ will have 0, 1 or 2 components. Let $n_\gamma^a(\beta)$ be this number and similarly define $n_\gamma^b(\beta)$. Note that if β is a closed leaf then the number will always be zero. The content of the following lemma is that the length function ℓ_γ is given by any of the following two linear

functions

$$\begin{aligned}\ell_\gamma^a : \mathbb{R}^{|\lambda|} &\rightarrow \mathbb{R}, \quad \ell_\gamma^a(\mathbf{x}) = \sigma_\gamma^a \sum_{i=1}^{|\lambda|} n_\gamma^a(\gamma_i) x_i \\ \ell_\gamma^b : \mathbb{R}^{|\lambda|} &\rightarrow \mathbb{R}, \quad \ell_\gamma^b(\mathbf{x}) = \sigma_\gamma^b \sum_{i=1}^{|\lambda|} n_\gamma^b(\gamma_i) x_i\end{aligned}$$

Lemma 3.2 *If γ is a closed curve in λ and $\ell_\gamma : \mathcal{T}(X) \rightarrow \mathbb{R}$ is its length function then*

$$\ell_\gamma = \ell_\gamma^a \circ s_\lambda = \ell_\gamma^b \circ s_\lambda.$$

Proof. Let (Y, f) be a marked hyperbolic structure in $\mathcal{T}(X)$. We will calculate $\ell_\gamma^a \circ s_\lambda(Y, f)$. Working in the universal cover $\mathbb{H}^2$ using the upper half space model we can assume that $\bar{f}(\tilde{\gamma})$ is the vertical line at $x = 0$.

If we assume that $\sigma_\gamma^a = -1$ then we can also choose $\tilde{f}$ so that the geodesics in $\tilde{f}(\tilde{\lambda})$ that intersect that a -side of $\tilde{f}(\tilde{\gamma})$ are vertical geodesics with negative x -coordinates. Label these x -coordinates x_i with $x_{i+1} < x_i$. By our normalization the subgroup of the deck group for Y that fixes $\tilde{f}(\tilde{\gamma})$ will be generated by the isometry $z \mapsto e^{\ell_\gamma(Y, f)} z$. The set of vertical geodesics will be invariant under this isometry and we will have $x_{i+k} = e^{\ell_\gamma(Y, f)} x_i$ where

$$k = \sum_{\beta \in \lambda} n_\gamma^a(\beta)$$

is the number of components of the intersection of λ with the a -side of γ . Therefore $\ell_\gamma(Y, f) = \log x_{i+k}/x_i$.

Let Δ_i be the ideal triangle that has two vertical sides with x -coordinate x_i and x_{i+1} . Note that the midpoints of the two vertical sides will have the same y -coordinate which we label m_i and that $x_{i+1} = x_i - m_i$. We also observe that $s(\Delta_i, \Delta_{i+1}) = -\log m_{i+1}/m_i$ so

$$\begin{aligned}\sum_{i=0}^{k-1} s(\Delta_i, \Delta_{i+1}) &= -\sum_{i=0}^{k-1} \log m_{i+1}/m_i \\ &= -\log m_k + \log m_0 \\ &= -\log(x_k - x_{k+1}) + \log(x_0 - x_1) \\ &= -\log x_k/x_0 \\ &= -\ell_\gamma(Y, f).\end{aligned}$$

The second to last equality follows from the fact that $\ell_\gamma(Y, f) = \log x_k/x_0 = \log x_{k+1}/x_1$ and therefore $x_{k+1}/x_k = x_1/x_0$.

To finish the proof we need to write the sum on the left in terms of our coordinates. To do so we note that each vertical geodesic with x -coordinate x_i maps to an isolated component β of λ and $s_\beta(Y, f) = s(\Delta_{i-1}, \Delta_i)$. Furthermore this geodesic will intersect the a -side of γ so $n_\gamma^a(\beta)$ is positive. In fact each component β of λ that intersects the a -side of γ will have exactly $n_\gamma^a(\beta)$ pre-images among the vertical geodesics with x -coordinates $x_1, \dots, x_k$. Therefore

$$\begin{aligned} \ell_\gamma^a \circ s_\lambda(Y, f) &= \sigma_\gamma^a \sum_{i=1}^{|\lambda|} n_\gamma^a(\gamma_i) s_{\gamma_i}(Y, f) \\ &= - \sum_{\beta \in \lambda} n_\gamma^a(\beta) s_\beta(Y, f) \\ &= - \sum_{i=0}^{k-1} s(\Delta_i, \Delta_{i+1}) \end{aligned}$$

which completes the proof when $\sigma_\gamma^a = -1$.

When $\sigma_\gamma^a = +1$ the proof is exactly the same except the vertical geodesics on the a -side of $\bar{f}(\tilde{\gamma})$ have positive x -coordinate. If we label the coordinates x_i with $x_i < x_{i+1}$ then $s(\Delta_i, \Delta_{i+1}) = \log m_{i+1}/m_i$. The rest of the proof of the proof is exactly the same so this accounts for the $\sigma_\gamma^a = +1$ in the definition of ℓ_γ^a .

The proof for ℓ_γ^b is obviously the same. 3.2

Now let c be a cusp of X . We let $n_c(\beta)$ be the number of components of the intersection of β with a horospherical neighborhood of c and define

$$\ell_c(\mathbf{x}) = \sum_{i=1}^{|\lambda|} n_c(\gamma_i) x_i.$$

We then have the following lemma.

Lemma 3.3 *If c is a cusp of X then $\ell_c \circ s_\lambda(Y, f) = 0$ for all (Y, f) in $\mathcal{T}(X)$.*

Proof. The proof follows the same basic idea as the proof of Lemma 3.2. We can assume that a component of the pre-image of the horosphere neighborhood of the cusp is a horosphere neighborhood of infinity in the upper half-space model of $\mathbb{H}^2$. Then the geodesics in $\bar{f}(\lambda)$ that intersect this neighborhood will be vertical geodesics with x -coordinates x_i labeled such that $x_i < x_{i+1}$. The neighborhood will be invariant under the deck transformation that fixes infinity and we can assume that it is of the form $z \mapsto z + 1$. In particular $x_{i+k} = x_i + 1$ where

$$k = \sum_{\beta \in \lambda} n_c(\beta).$$

The set of midpoints m_i will also be invariant under the action of $z \mapsto z + 1$ so we also have $m_i = m_{i+k}$. Repeating the calculations from Lemma 3.2 we see that

$$\sum_{i=0}^{k-1} s(\Delta_i, \Delta_{i+1}) = \log x_k/x_0 = 0$$

and that

$$\ell_c \circ s_\lambda(Y, f) = \sum_{i=0}^{k-1} s(\Delta_i, \Delta_{i+1})$$

and the lemma is proven. 3.3

The functions $\ell_\gamma^a, \ell_\gamma^b$ and ℓ_c are linear functions defined on all of $\mathbb{R}^{|\lambda|}$, in particular we deduce from Lemma 3.2 and Lemma 3.3:

Corollary 3.4 *The image of s_λ is contained in the convex polytope $T_\lambda \subset \mathbb{R}^{|\lambda|}$ consisting of all those $\mathbf{x}$ with $\ell_\gamma^a(\mathbf{x}) = \ell_\gamma^b(\mathbf{x}) > 0$ for all simple closed curves γ in λ and $\ell_c(\mathbf{x}) = 0$ for all cusps of X .* 3.4

3.4 The coordinates are coordinates

We prove now the central result of this section:

Theorem 3.5 *The map s_λ is a homeomorphism from $\mathcal{T}(X)$ to T_λ .*

We will derive Theorem 3.5 from some standard results about Teichmüller space and the following proposition.

Proposition 3.6 *Let α be a closed curve on X and $\ell_\alpha : \mathcal{T}(X) \rightarrow \mathbb{R}^+$ its length function. Then there is a convex function $\bar{\ell}_\alpha : \mathbb{R}^{|\lambda|} \rightarrow \mathbb{R}$ such that $\ell_\alpha = \bar{\ell}_\alpha \circ s_\lambda$. Furthermore the $\bar{\ell}_\alpha$ -image of T_λ is in $\mathbb{R}^+$ and if α intersects every leaf of λ then $\bar{\ell}_\alpha$ is strictly convex.*

Proposition 3.6 is the main work of this paper; we defer its proof to the final section.

Proof of Theorem 3.5. As mentioned before, the fact that s_λ can be expressed in terms of cross ratios shows that it is continuous. By Corollary 3.4 the image lies in T_λ which is a convex subset of a linear subspace of $\mathbb{R}^{|\lambda|}$. The number of ideal triangles in $X \setminus \lambda$ is $-2\chi(X) = \text{area}(X)/\pi$ and therefore $|\lambda| = -3\chi(X) + n_c$ where n_c is the number of closed curves in λ . In the definition of T_λ there is one equation for each closed curve and one equation for each cusp. These equations are also linearly independent so T_λ is

an open subset of a linear subspace of dimension $|\lambda| - n_c - n_p$ where n_p is the number of punctures (or cusps). Note that $\dim \mathcal{T}(X) = -3\chi(X) - n_p = |\lambda| - n_c - n_p$.

If $s_\lambda(Y_0) = s_\lambda(Y_1)$ then by Proposition 3.6, $\ell_\alpha(Y_0) = \ell_\alpha(Y_1)$ for all closed curves α and therefore $Y_0 = Y_1$. This implies that s_λ is injective. By invariance of domain s_λ is a local homeomorphism and its image an open subset of T_λ .

To see that the image is closed take a finite collection of closed curves $\Gamma = \{\alpha_1, \dots, \alpha_k\}$ such that the complementary pieces are disks or punctured disks. Such a collection *binds* the surface. Let $\ell_\Gamma : \mathcal{T}(X) \rightarrow \mathbb{R}^+$ be the sum of the length functions ℓ_{α_i} and similarly define $\bar{\ell}_\Gamma : \mathbb{R}^{|\lambda|} \rightarrow \mathbb{R}$ and the sum of the $\bar{\ell}_{\alpha_i}$.

Let Y_i be a sequence in $\mathcal{T}(X)$ such that $s_\lambda(Y_i)$ converges to a point $\mathbf{x}$ in T_λ . By Proposition 3.6 $\bar{\ell}_\Gamma(\mathbf{x}) < \infty$ and since $\ell_\Gamma(Y_i) = \bar{\ell}_\Gamma \circ s_\lambda(Y_i) \rightarrow \bar{\ell}_\Gamma(\mathbf{x})$ we see that $\ell_\Gamma(Y_i)$ is uniformly bounded. Therefore the Y_i lie in a compact subset of $\mathcal{T}(X)$ by Lemma 3.1 of [7] or Proposition 2.4 of [9]. In particular we can find a subsequence such that Y_{i_k} converges to some $Y \in \mathcal{T}(X)$ and therefore $\lim s_\lambda(Y_i) = \lim s_\lambda(Y_{i_k}) = s_\lambda(Y)$ is in image of s_λ . Therefore the image is a closed set.

Since the image is open and closed it must be all of T_λ . 3.5

Before moving observe that Theorem 1.3 is an immediate consequence of Proposition 3.6 and Theorem 3.5:

Theorem 1.3 *Let X be a complete hyperbolic surface with finite area of genus g and with n cusps. Let λ be a maximal lamination in X with finitely many leaves and let $s_\lambda : \mathcal{T}(X) \xrightarrow{\sim} T_\lambda$ be the shearing coordinates associates to λ . For any essential curve γ in X the function*

$$l_\gamma \circ s_\lambda^{-1} : T_\lambda \rightarrow \mathbb{R}_+$$

is convex. If moreover the curve γ intersects all the leaves of λ then $l_\gamma \circ s_\lambda^{-1}$ is strictly convex. 1.3

3.5 The Nielsen realization conjecture

Kerckhoff proved the follow theorem using the convexity of length functions along earthquake paths. We give a similar proof using our convexity result.

Theorem 3.7 (Kerckhoff) *If Γ binds then ℓ_Γ has a unique minimum on $\mathcal{T}(X)$.*

Proof. Fix an ideal triangulation λ . Since Γ binds there must be some curve in Γ that is not in λ so by Proposition 3.6 the length function $\bar{\ell}_\Gamma$ is strictly convex on T_λ . As we noted in the proof of Theorem 3.5 the function ℓ_Γ is proper and since $\ell_\Gamma = \bar{\ell}_\Gamma \circ s_\lambda$ the function $\bar{\ell}_\Gamma$ is also proper. A proper, strictly convex function that is bounded below has a unique minimum so $\bar{\ell}_\Gamma$, and therefore ℓ_Γ , have unique minimums. 3.7

Kerckhoff used this result to prove the Nielsen realization conjecture. As the proof is short we include it here.

Theorem 1.2 (Kerckhoff) *The action of every finite subgroup of the mapping class group on $\mathcal{T}(S)$ has a fixed point.*

Proof. Let G be the finite subgroup. The G orbit of any finite set of binding curves will still bind and will be G -invariant. Let Γ be such a G -invariant binding set and let X be the unique minimum of ℓ_Γ . Clearly X is fixed by G . 1.2

3.6 Surfaces with geodesic boundary

We conclude this section with a few remarks on surfaces with boundary. Let X be a finite area hyperbolic surface with boundary components $\beta_1, \dots, \beta_k$. For each boundary component β_i choose an interval I_i of $(0, \infty)$ and let $\mathcal{T}(X; I_1, \dots, I_k)$ be the Teichmüller space of marked hyperbolic surfaces where the length of β_i is in the interval I_i . We allow the I_i to be open, closed, half-open or a point.

An ideal triangulation of X is still a finite leaved geodesic lamination whose complement is the union of an open ideal triangle; in particular, $\partial X \subset \lambda$. Let $\hat{\lambda} = \lambda \setminus \partial X$ be the union of the all interior leaves of λ . The definition of the coordinate map $s_\lambda : \mathcal{T}(X; I_1, \dots, I_k) \rightarrow \mathbb{R}^{|\hat{\lambda}|}$ still makes sense. For each boundary component β of X we have a function $\ell_\beta : \mathbb{R}^{|\hat{\lambda}|} \rightarrow \mathbb{R}$ where the composition $\ell_\beta \circ s_\lambda$ is the length function for the boundary component. The definition of ℓ_β is exactly the same as the definition of the functions ℓ_γ^a and ℓ_γ^b . Let $T_{\lambda, I_1, \dots, I_k}$ be the subset of $T_\lambda \subset \mathbb{R}^{|\hat{\lambda}|}$ satisfying $\ell_{\beta_i}(\mathbf{x}) \in I_i$ for $i = 1, \dots, k$.

Theorem 3.8 *The map s_λ is a homeomorphism from $\mathcal{T}(X; I_1, \dots, I_k)$ to $T_{\lambda, I_1, \dots, I_k}$.*

Proof. Note that if each I_i is a point then T_λ is an open subset of a linear subspace of $\mathbb{R}^{|\hat{\lambda}|}$ and the proof is the same as the proof Theorem 3.5. The general case follows from this observation. 3.8

4 Fenchel-Nielsen coordinates

In this section we prove Theorem 1.1. We will assume knowledge of some basic form of the Fenchel-Nielsen coordinates; see for example [2].

Let X be a hyperbolic surface and γ a non-peripheral simple closed curve. Fenchel and Nielsen defined for $t \in \mathbb{R}$ the twist deformation $T_\gamma^t(X)$ of X along γ with twist

parameter t as follows. First isotope γ to the geodesic in its free homotopy class, also denoted by γ , and let $g : \mathbb{R} \rightarrow \gamma$ be a parametrization by arc-length. Let γ_1 and γ_2 be the two boundary components obtained after cutting X along γ and for $i = 1, 2$ let $g_i : \mathbb{R} \rightarrow \gamma_i$ be the induced parametrization. Up to relabeling, we may assume that g_1 is orientation preserving and g_2 is orientation reversing with respect to the induced orientation of the boundary curves γ_1 and γ_2 . The hyperbolic surface $T_\gamma^t(X)$ is obtained from the cut open surface by identifying the points $g_1(s)$ with $g_2(s + t)$ for all $s \in \mathbb{R}$. Compare with figure 5.




Figure 5: Twist deformation for t equal to the length of the bold printed arc

The surface $T_\gamma^t(X)$ is just a hyperbolic surface; in particular it has so far no marking. However, it is well-known that there is flow, the Fenchel-Nielsen Dehn-twist flow

$$\tau_\gamma : \mathbb{R} \times \mathcal{T}(X) \rightarrow \mathcal{T}(X), \quad (t, (Y, f)) \mapsto \tau_\gamma^t(Y, f)$$

such that for $(Y, f) \in \mathcal{T}(X)$, the surface $T_\gamma^t(Y)$ is the hyperbolic surface associated to the point $\tau_\gamma^t(Y, f)$.

If γ and γ' are disjoint curves, then the flows τ_γ and $\tau_{\gamma'}$ commute. In particular, labeling by $\gamma_1, \dots, \gamma_{|\mathcal{P}|}$ the components of some pants decomposition $\mathcal{P}$ of X we have an action

$$\tau_{\mathcal{P}} : \mathbb{R}^{|\mathcal{P}|} \times \mathcal{T}(X) \rightarrow \mathcal{T}(X), \quad ((t_1, \dots, t_{|\mathcal{P}|}), (Y, f)) \mapsto \tau_{\mathcal{P}}^{(t_1, \dots, t_{|\mathcal{P}|})}(Y, f)$$

where $\tau_{\mathcal{P}}^{(t_1, \dots, t_{|\mathcal{P}|})} = \tau_{\gamma_1}^{t_1} \circ \dots \circ \tau_{\gamma_{|\mathcal{P}|}}^{t_{|\mathcal{P}|}}$. It is well-known that the action given $\tau_{\mathcal{P}}$ is free and proper.

To the pants decomposition $\mathcal{P}$ of X we can also associate the function

$$\ell_{\mathcal{P}} : \mathcal{T}(X) \longrightarrow (\mathbb{R}^+)^{|\mathcal{P}|}$$

which assigns to each point in $\mathcal{T}(X)$ the $|\mathcal{P}|$ -tuple of lengths of curves in the pants decomposition. Observe that the $\mathbb{R}^{|\mathcal{P}|}$ -action $\tau_{\mathcal{P}}$ preserves by definition the function $\ell_{\mathcal{P}}$ and hence acts on the fibers.

In fact, any existence results for Fenchel-Nielsen coordinates implies that the $\tau_{\mathcal{P}}$ -orbits actually coincide with the fibers of $\ell_{\mathcal{P}}$. We can summarize this discussion as follows:

Proposition 4.1 *Let $\mathcal{P}$ be a pants decomposition of a finite area surface X . Then*

$$\ell_{\mathcal{P}} : \mathcal{T}(X) \rightarrow (\mathbb{R}^+)^{|\mathcal{P}|}$$

has a natural structure as a $\mathbb{R}^{|\mathcal{P}|}$ -principal bundle.

Every principal bundle over a contractible space is trivial but not canonically trivialized. In fact, any choice of a section yields a trivialization and vice-versa. From the point of view of the authors, all coordinates for Teichmüller space obtained by trivializing the principal bundle $\ell_{\mathcal{P}} : \mathcal{T}(X) \rightarrow (\mathbb{R}^+)^{|\mathcal{P}|}$ deserve to be referred to as *Fenchel-Nielsen coordinates*. We are now ready to prove Theorem 1.1.

Theorem 1.1. *Let X be a complete, finite area, hyperbolic surface of genus g and with n cusps, and fix a pants decomposition $\mathcal{P}$ of X . There are Fenchel-Nielsen coordinates $\Phi : \mathcal{T}(X) \xrightarrow{\sim} \mathbb{R}_+^{3g+n-3} \times \mathbb{R}^{3g+n-3}$ associated to $\mathcal{P}$ such that for any essential curve γ in X the function*

$$l_{\gamma} \circ \Phi^{-1} : \mathbb{R}_+^{3g+n-3} \times \mathbb{R}^{3g+n-3} \rightarrow \mathbb{R}_+$$

is convex. If moreover the curve γ intersects all the components of $\mathcal{P}$ then $l_{\gamma} \circ \Phi$ is strictly convex.

To begin with extend $\mathcal{P}$ to an ideal triangulation λ and let

$$s_{\lambda} : \mathcal{T}(X) \rightarrow T_{\lambda}$$

be the shearing coordinates associated to λ . Theorem 1.1 will follow immediately from Theorem 1.3 when we interpret the shearing coordinates s_{λ} as Fenchel-Nielsen coordinates.

Denote by $\lambda_0 = \lambda \setminus \mathcal{P}$ the set of isolated leaves in λ , recall the definition of the convex polytope $T_{\lambda} \subset \mathbb{R}^{|\lambda|} = \mathbb{R}^{|\mathcal{P}|} \times \mathbb{R}^{|\lambda_0|}$ and observe that the factor $\mathbb{R}^{|\mathcal{P}|} \times \{0\}$ in the above splitting of $\mathbb{R}^{\lambda}$ is contained in T_{λ} . In particular, the canonical $\mathbb{R}^{|\mathcal{P}|}$ -principal bundle structure on

$$\pi : \mathbb{R}^{|\lambda|} = \mathbb{R}^{|\mathcal{P}|} \times \mathbb{R}^{|\lambda_0|} \rightarrow \mathbb{R}^{|\lambda_0|}$$

induces a $\mathbb{R}^{|\mathcal{P}|}$ -principal bundle structure on

$$\pi : T_{\lambda} \rightarrow \pi(T_{\lambda})$$

Here π is the projection to the second factor of the splitting of $\mathbb{R}^{|\lambda|}$.

It follows directly from the definition just before Lemma 3.2 that for all $\gamma \in \mathcal{P}$ the linear form $\ell_\gamma^a : T_\lambda \rightarrow \mathbb{R}^+$ is independent of the $\mathbb{R}^{|\mathcal{P}|}$ -factor and hence induces a well-defined linear function $\hat{\ell}_\gamma$ on $\pi(T_\lambda)$. Observe that this function is positive by Lemma 3.2. Denoting by $\hat{\ell}_\mathcal{P} : \pi(T_\lambda) \rightarrow (\mathbb{R}^+)^{|\mathcal{P}|}$ the function whose γ -coordinate is $\hat{\ell}_\gamma$ we have from Lemma 3.2 that the following diagram commutes

$$\begin{array}{ccc} \mathcal{T}(X) & \xrightarrow{s_\lambda} & T_\lambda \\ \ell_\mathcal{P} \downarrow & & \downarrow \pi \\ (\mathbb{R}^+)^{|\mathcal{P}|} & \xleftarrow{\hat{\ell}_\mathcal{P}} & \pi(T_\lambda) \end{array} \quad (4.3)$$

The map $\hat{\ell}_\mathcal{P}$ is then linear and surjective. Since

$$\dim(\pi(T_\lambda)) \leq \dim(T_\lambda) - \dim(\mathbb{R}^{|\mathcal{P}|}) \leq |\mathcal{P}|$$

we obtain that $\hat{\ell}_\mathcal{P}$ is a linear isomorphism. In particular, the bundles $\mathcal{T}(X) \rightarrow (\mathbb{R}^+)^{|\mathcal{P}|}$ and $T_\lambda \rightarrow \pi(T_\lambda)$ are isomorphic as fiber bundles. Moreover, it follows directly from the definition of the shearing coordinates that they are also equivalent as $\mathbb{R}^{|\mathcal{P}|}$ -principal bundles. Compare with figure 6.

The projection $T_\lambda \rightarrow (\mathbb{R}^+)^{|\mathcal{P}|}$ is linear and has kernel of dimension $|\mathcal{P}|$. In particular, there is a linear map $L : T_\lambda \rightarrow \mathbb{R}^{|\mathcal{P}|} \times (\mathbb{R}^+)^{|\mathcal{P}|}$ such that the following diagram commutes:

$$\begin{array}{ccc} T_\lambda & \xrightarrow{L} & \mathbb{R}^{|\mathcal{P}|} \times (\mathbb{R}^+)^{|\mathcal{P}|} \\ & \searrow \pi & \swarrow \\ & (\mathbb{R}^+)^{|\mathcal{P}|} & \end{array} \quad (4.4)$$

Here the unlabeled arrow is the projection on the first factor.

Combining (4.3) and (4.4) we obtain a trivialization of the principal bundle $\ell_\mathcal{P} : \mathcal{T}(X) \rightarrow (\mathbb{R}^+)^{|\mathcal{P}|}$, i.e. Fenchel-Nielsen coordinates, which differ from the shearing coordinates s_λ by a linear map. Since by Theorem 1.3 the length functions are convex with respect to the shearing coordinates and convexity is preserved by linear maps, the same result holds for this choice of Fenchel-Nielsen coordinates. This concludes the proof of Theorem 1.1. 1.1

5 Length functions on $\mathcal{T}(A)$

Let A be a complete hyperbolic annulus and assume that the core curve α is isotopic to a geodesic and let $\mathcal{T}(A)$ be the Teichmüller space of A . Observe that according to the




Figure 6: The lower horizontal arrow is the twist deformation along the central curve γ where the parameter is the length of the bold printed arc. The pictures in the upper row schematize the situation in the universal cover: the straight lines represent a lift $\tilde{\gamma}$ of γ , the white dots are the midpoints of the sides of the chosen triangles Δ^a and Δ^b ; the black dots are the projections of these midpoints to $\tilde{\gamma}$. The increase in the shearing coordinate associated to γ coincides with the twist parameter.

definition above, $\mathcal{T}(A)$ is infinite dimensional. Fix also an ideal triangulation $\tilde{\lambda}$ of A and let λ be the sublamination of $\tilde{\lambda}$ obtained by deleting all leaves of λ which are disjoint from the core geodesic of A . Assume that λ contains only finitely many non-isolated leaves. In this section we construct a map

$$s_\lambda : \mathcal{T}(A) \rightarrow \mathbb{R}^{|\lambda|}$$

and a convex function

$$\bar{\ell} : \mathbb{R}^{|\lambda|} \longrightarrow (0, \infty]$$

such that $\ell = \bar{\ell} \circ s_\lambda$. Here

$$\ell : \mathcal{T}(A) \longrightarrow (0, \infty)$$

is the function which assigns to each hyperbolic annulus the length of its core curve.

5.1 A treatise on wedges

An *ideal wedge* is the region bounded by two asymptotic geodesics in $\mathbb{H}^2$. If W is an ideal wedge in $\mathbb{H}^2$ or A then there is a unique ideal triangle $\Delta(W)$ two of whose boundary components are the boundary components of W . In particular, the two boundary

edges γ_1 and γ_2 of the ideal wedge W have well-determined *midpoints*. Orienting both boundary edges we parametrize γ_1 and γ_2 by $\mathbb{R}$ via the signed distance from their. To $(x, y) \in \mathbb{R}^2$ we associate the pair of points in ∂W corresponding to x in γ_1 and y in γ_2 respectively; let $d(x, y)$ be the distance in $\mathbb{H}^2$ of these two points in ∂W .




Figure 7: The length of the bold segment is by definition $d(x, y)$.

We can describe the function

$$d : \mathbb{R}^2 \longrightarrow (0, \infty)$$

more explicitly using the upper half space model of $\mathbb{H}^2$. Namely, define $d(x, y)$ to be the distance in $\mathbb{H}^2$ between the points ie^x and $1 + ie^y$.

Lemma 5.1 *The function d is strictly convex and*

$$|x - y| \leq d(x, y) \leq |x - y| + \frac{1}{\max\{e^x, e^y\}}.$$

Lemma 5.1 can be proved via a simple computation but it should be remarked that in more intrinsic terms it just follows from the fact that $\mathbb{H}^2$ is negatively curved and hence that the distance function is convex. 5.1

An injective isometric immersion of a wedge into a hyperbolic annulus A will also be called an ideal wedge; all the definitions above carry over without difficulties.

An *ideal wedgelation* is a geodesic lamination λ on A whose complement is a disjoint union of ideal wedges, and such that λ contains only finitely many non-isolated leaves. For the sake of concreteness we will also assume that every leaf of λ which is isolated to one side is actually isolated.

Lemma 5.2 *Let S be a hyperbolic surface and λ an ideal triangulation of S . Let also A be an annulus and $\pi : A \rightarrow S$ a covering. The set of leaves of $\pi^{-1}(\lambda)$ which intersect every curve in A homotopic to the core curve is an ideal wedgelation. 5.2*

On a finite area surface it is not possible to consistently orient an ideal triangulation. On an annulus, an ideal wedgelation can be consistently oriented and this will be important in the work below. To do so we fix an orientation on A and of its core geodesic α . Observe that all the leaves of an ideal wedgelation λ of A intersect α exactly once. We then orient every geodesic in λ so that if γ is a geodesic in λ the orientation of A at $\alpha \cap \gamma$ is given by the ordered pair of the oriented tangent vectors to α and γ .




Figure 8: An ideal wedgelation of the annulus A with the corresponding orientations induced by the bold printed core geodesic.

Observe that if W is an ideal wedge of in $A \setminus \lambda$, then the orientations of the boundary edges of W determined by W and the orientations determined by λ agree if and only if the vertex of W is to the left of the core curve α .

5.2 The shearing map

Recall that for every ideal wedge W in $\mathbb{H}^2$ or A we have associated a unique ideal triangle $\Delta(W)$. If W^a and W^b are disjoint wedges in $\mathbb{H}^2$ and γ is a geodesic in $\mathbb{H}^2$ separating W^a from W^b and asymptotic to some boundary edge of W^a and some boundary edge W^b we define

$$s(W^a, W^b, \gamma) = s(\Delta(W^a), \Delta(W^b), \gamma).$$

Clearly, all claims of Lemma 3.1 hold in this setting as well.

An ideal wedgelation of A yields now a map

$$s_\lambda : \mathcal{T}(A) \longrightarrow \mathbb{R}^{|\lambda|}$$

in the same way as it did for an ideal triangulation on a surface except we replace ideal triangles with ideal wedges W_γ^- and W_γ^+ . We remark that s_λ does not yield coordinates of $\mathcal{T}(A)$. We refer to the map s_λ as the *shearing map*.

5.3 The function L

Let λ be an ideal wedgelation of A , $\lambda_0 \subset \lambda$ the collection of isolated leaves of λ and $\mathcal{W}$ the set of ideal wedges in $A \setminus \lambda$. Recall that $\lambda \setminus \lambda_0$ consists by definition of only finitely many leaves and that we are also working under the additional assumption that the leaves in $\lambda \setminus \lambda_0$ are non-isolated on both sides.

For each $W \in \mathcal{W}$ let γ_W^- be the left boundary edge and γ_W^+ be the right boundary edge where left and right is defined with respect to the orientation of the core curve α . Observe that both γ_W^+ and γ_W^- belong to λ . If the two boundary components γ_W^+ and γ_W^- of W are asymptotic to the left of α then define $d_W(x, y) = d(x, y)$ for $(x, y) \in \mathbb{R}^2$. If they are asymptotic to the right of α then define $d_W(x, y) = d(-x, -y)$. Here left and right are defined with respect to the orientation of α .

Define a function $L : \mathbb{R}^{|\lambda|} \times \mathbb{R}^{|\lambda_0|} \longrightarrow (0, \infty]$ by

$$L(\mathbf{x}, \mathbf{y}) = \sum_{W \in \mathcal{W}} d_W(\gamma_W^-(\mathbf{y}), \gamma_W^+(\mathbf{y}) + \gamma_W^+(\mathbf{x})).$$

Here $\gamma(\mathbf{x})$ is the γ -coordinate of $\mathbf{x} \in \mathbb{R}^\lambda$. Note that L does not depend on the coordinates of the non-isolated leaves in λ .

Meaning of L : The definition of L is a bit obscure. Before moving on we explain its meaning in the particular case that λ is a finite wedgelation. In particular $\lambda_0 = \lambda$ and the sum in the definition of L is finite. Given a point $(B, f) \in \mathcal{T}(A)$, its image $\mathbf{x} = s_\lambda(B, f)$ under the shearing map encodes how the different wedges are glued in X . The element $\mathbf{y} \in \mathbb{R}^{\lambda_0}$ picks a point in each leaf of the (finite) wedgelation λ . In particular, $\mathbf{y}$ picks in every wedge W two different points in the boundary. Let α_W be the geodesic segment in W joining these two points. The juxtaposition of all the segments α_W is a closed loop in (B, f) homotopic to the core curve; $L(\mathbf{x}, \mathbf{y})$ is the length of this loop.

5.4 Piecewise geodesic segments

Our next aim is to generalize the preceding discussion on the meaning of the function L . Assume now that λ is a general wedgelation and recall that λ_0 is the collection of isolated leaves of λ .

For a marked hyperbolic annulus $(B, f) \in \mathcal{T}(A)$ let $\mathcal{P}(B, f)$ be the set of closed continuous loops β on B homotopic to $f(\alpha)$ and such that the intersection of β with

each wedge in $B \setminus \bar{f}(\lambda)$ is a geodesic segment. In particular, any $\beta \in \mathcal{P}(B, f)$ intersects each geodesic in $\bar{f}(\lambda)$ exactly once. We let

$$p_\lambda : \mathcal{P}(B, f) \longrightarrow \mathbb{R}^{|\lambda_0|}$$

be the map defined by the property that $\gamma(p_\lambda(\beta))$ is the signed distance between $m_{\bar{f}(\gamma)}$ and the intersection of β with $\bar{f}(\gamma)$ for all $\gamma \in \lambda_0$. As always, the sign is determined by the orientation induced by the core geodesic α .

The map p_λ is clearly injective. However, if $\lambda_0 \neq \lambda$ it is easy to see that it is not surjective. Our next goal is to determine the image of p_λ . In order to do so we define two quantities $a_\gamma^-(\mathbf{x}, \mathbf{y})$ and $a_\gamma^+(\mathbf{x}, \mathbf{y})$ for every non-isolated leaf γ ; recall that we are assuming that leaves which are isolated to one side are actually isolated. We first define a_γ^- . Label the geodesics between γ and W_γ^- , the (arbitrarily) chosen wedge to the left of γ , by $\gamma_0^- = \gamma_{W_\gamma^-}^+, \gamma_1^-, \dots$ so that they are positively ordered with respect to α . Define

$$a_\gamma^-(\mathbf{x}, \mathbf{y}) = \lim_{n \rightarrow \infty} \left(\gamma_n^-(\mathbf{y}) + \sum_{i=0}^n \gamma_i^-(\mathbf{x}) \right) \quad (5.5)$$

if the limit exists. The definition of $a_\gamma^+(\mathbf{x}, \mathbf{y})$ is very similar with small changes. Label the geodesics between W_γ^+ and γ by $\gamma_0^+ = \gamma_{W_\gamma^+}^-, \gamma_1^+, \dots$ so that they are negatively ordered with respect to γ and define

$$a_\gamma^+(\mathbf{x}, \mathbf{y}) = \lim_{n \rightarrow \infty} \left(\gamma_n^+(\mathbf{y}) - \sum_{i=0}^{n-1} \gamma_i^+(\mathbf{x}) \right) \quad (5.6)$$

if the limit exist and $a_\gamma^\pm(x, y) = \infty$ otherwise. To understand the meaning of the sums in the definition compare with the last claim in Lemma 3.1.

Given $\mathbf{x} \in \mathbb{R}^{|\lambda|}$ define $P_\mathbf{x} \subset \mathbb{R}^{|\lambda_0|}$ to be the set

$$P_\mathbf{x} = \left\{ \mathbf{y} \in \mathbb{R}^{|\lambda_0|} \mid \begin{array}{l} \text{the limits (5.5) and (5.6) exist, and} \\ a_\gamma^-(\mathbf{x}, \mathbf{y}) = a_\gamma^+(\mathbf{x}, \mathbf{y}) + \gamma(\mathbf{x}) < \infty \text{ for all } \gamma \in \lambda \setminus \lambda_0 \end{array} \right\}.$$

We prove:

Lemma 5.3 *Let (B, f) be a marked hyperbolic annulus in $\mathcal{T}(A)$. If $\mathbf{x} = s_\lambda(B, f)$ then the p_λ -image of $\mathcal{P}(B, f)$ is $P_\mathbf{x}$ and*

$$\text{length}(\beta) = L(\mathbf{x}, p_\lambda(\beta)).$$

Proof. Let γ be a geodesic in $\lambda \setminus \lambda_0$. As above we label the geodesics on the left of γ by $\gamma_0^-, \gamma_1^-, \dots$ and on the right by $\gamma_0^+, \gamma_1^+, \dots$. If β is a curve in $\mathcal{P}(B, f)$ let $p_i^\pm$ be the point of intersection of β with $\bar{f}(\gamma_i^\pm)$. Note that the sequence p_i^- and p_i^+ will limit to the same point p on γ since β is a continuous path. Therefore both $h_{\bar{f}(\gamma_i^-), \bar{f}(\gamma)}(p_i^-)$ and $h_{\bar{f}(\gamma_i^+), \bar{f}(\gamma)}(p_i^+)$ will limit to p . Let d_i^- be the signed distance between $h_{\bar{f}(\gamma_0^-), \bar{f}(\gamma)}(m_{\bar{f}(\gamma_0^-)})$ and $h_{\bar{f}(\gamma_i^-), \bar{f}(\gamma)}(p_i^-)$ and similarly define d_i^+ . Then d_i^- will limit to the signed distance between $h_{\bar{f}(\gamma_0^-), \bar{f}(\gamma)}(m_{\bar{f}(\gamma_0^-)})$ and p which we label d^- . Similarly d^+ , the limit of d_i^+ , will be the signed distance between $h_{\bar{f}(\gamma_0^+), \bar{f}(\gamma)}(m_{\bar{f}(\gamma_0^+)})$ and p . Since the signed distance between $h_{\bar{f}(\gamma_0^-), \bar{f}(\gamma)}(m_{\bar{f}(\gamma_0^-)})$ and $h_{\bar{f}(\gamma_0^+), \bar{f}(\gamma)}(m_{\bar{f}(\gamma_0^+)})$ is $\gamma(\mathbf{x})$ we have

$$d^- - d^+ = \gamma(\mathbf{x}). \quad (5.7)$$

Since the h -maps are isometries and $h_{\bar{f}(\gamma_i^-), \bar{f}(\gamma)} \circ h_{\bar{f}(\gamma_0^-), \bar{f}(\gamma_i^-)} = h_{\bar{f}(\gamma_0^-), \bar{f}(\gamma)}$ we have that d_i^- is also equal to the signed distance between $h_{\bar{f}(\gamma_0^-), \bar{f}(\gamma)}(m_{\bar{f}(\gamma_0^-)})$ and p_i^- . By definition $\gamma_n^-(p_\lambda(\beta))$ is the signed distance between $m_{\bar{f}(\gamma_n^-)}$ and p_n^- so we have

$$d_n^- = \gamma_n^-(p_\lambda(\beta)) + \sum_{i=0}^n \gamma_i^-(\mathbf{x})$$

by the last claim of Lemma 3.1. Therefore $d^- = a_\gamma^-(\mathbf{x}, p_\lambda(\beta))$ and similarly $d^+ = a_\gamma^+(\mathbf{x}, p_\lambda(\beta))$. Rearranging (5.7) we have

$$a_\gamma^-(\mathbf{x}, p_\lambda(\beta)) = a_\gamma^+(\mathbf{x}, p_\lambda(\beta)) + \gamma(\mathbf{x})$$

and therefore $p_\lambda(\beta)$ is in $P_\mathbf{x}$.

For every $\mathbf{y} \in P_\mathbf{x}$ we need to build a curve β on B such that $p_\lambda(\beta) = \mathbf{y}$. Given a geodesic α in λ_0 let p_α be the point on $\bar{f}(\alpha)$ whose signed distance from $m_{\bar{f}(\alpha)}$ is $\alpha(\mathbf{y})$. Then on each of the finitely many components of $B \setminus \bar{f}(\lambda \setminus \lambda_0)$ there is an arc that intersects each α in λ_0 at p_α and is a geodesic in each wedge. For these arcs to complete to a simple closed curve in $\mathcal{P}(B, f)$ we need that for each geodesic γ in $\lambda \setminus \lambda_0$ the limit of the points $p_{\gamma_i^-}$ is equal to the limit of $p_{\gamma_i^+}$. From the first paragraph of this proof we see that this holds when $\mathbf{y}$ is in $P_\mathbf{x}$.

Finally we note that the length of β is the sum of the lengths of the restriction of β to each ideal wedge. This is exactly the sum $L(\mathbf{x}, p_\lambda(\beta))$. 5.3

Before moving on to more interesting topics, we observe:

Lemma 5.4 *The set $\mathbb{P} = \{(\mathbf{x}, \mathbf{y}) \in \mathbb{R}^{|\lambda|} \times \mathbb{R}^{|\lambda_0|} | \mathbf{y} \in P_\mathbf{x}\}$ is a linear subspace and the projection $\mathbb{P} \rightarrow \mathbb{R}^{|\lambda|}$ is surjective.* 5.4

5.5 Convexity

We restrict from now on the function L to the linear subspace $\mathbb{P}$. The function L is an infinite sum of convex functions. This sum will not be finite everywhere so we need an extended notion of convexity. If V is vector space and $f : V \rightarrow (-\infty, \infty]$ is a function then f is *convex* if

$$f\left(\frac{\mathbf{x}_0 + \mathbf{x}_1}{2}\right) \leq \frac{f(\mathbf{x}_0) + f(\mathbf{x}_1)}{2}$$

whenever $f(\mathbf{x}_0)$ and $f(\mathbf{x}_1)$ are finite. The function is *strictly convex* if the inequality is always strict. We need the following general lemma.

Lemma 5.5 *Let V_0 and V_1 be vector spaces and P a subspace of $V_0 \times V_1$ whose projection onto V_0 is onto. Let $P_{\mathbf{x}} = \{\mathbf{y} \in V_1 | (\mathbf{x}, \mathbf{y}) \in P\}$. If $F : V_0 \times V_1 \rightarrow (0, \infty]$ is convex then*

$$f(\mathbf{x}) = \inf_{\mathbf{y} \in P_{\mathbf{x}}} F(\mathbf{x}, \mathbf{y})$$

is convex. If

$$f(\mathbf{x}) = \min_{\mathbf{y} \in P_{\mathbf{x}}} F(\mathbf{x}, \mathbf{y})$$

and F is strictly convex on P then f is strictly convex.

Proof. Let $\mathbf{x}_0$ and $\mathbf{x}_1$ be points in V_0 . We can assume that both $f(\mathbf{x}_0)$ and $f(\mathbf{x}_1)$ are finite for otherwise the lemma is trivial. For any $\epsilon > 0$ we can choose a $\mathbf{y}_0 \in P_{\mathbf{x}_0}$ and $\mathbf{y}_1 \in P_{\mathbf{x}_1}$ such that $f(\mathbf{x}_i) > F(\mathbf{x}_i, \mathbf{y}_i) - \epsilon$. Since P is a subspace we have that $(\mathbf{y}_0 + \mathbf{y}_1)/2$ is in $P_{(\mathbf{x}_0 + \mathbf{x}_1)/2}$ and therefore

$$\begin{aligned} f\left(\frac{\mathbf{x}_0 + \mathbf{x}_1}{2}\right) &\leq F\left(\frac{\mathbf{x}_0 + \mathbf{x}_1}{2}, \frac{\mathbf{y}_0 + \mathbf{y}_1}{2}\right) \\ &\leq \frac{F(\mathbf{x}_0, \mathbf{y}_0) + F(\mathbf{x}_1, \mathbf{y}_1)}{2} \\ &\leq \frac{f(\mathbf{x}_0) + f(\mathbf{x}_1)}{2} + \epsilon. \end{aligned}$$

Since the ϵ is arbitrary we have that

$$f\left(\frac{\mathbf{x}_0 + \mathbf{x}_1}{2}\right) \leq \frac{f(\mathbf{x}_0) + f(\mathbf{x}_1)}{2}$$

and f is convex.

If f is defined by a minimum then there are $\mathbf{y}_0$ and $\mathbf{y}_1$ such that $f(\mathbf{x}_i) = F(\mathbf{x}_i, \mathbf{y}_i)$. If F is also strictly convex we have

$$\begin{aligned} f\left(\frac{\mathbf{x}_0 + \mathbf{x}_1}{2}\right) &\leq F\left(\frac{\mathbf{x}_0 + \mathbf{x}_1}{2}, \frac{\mathbf{y}_0 + \mathbf{y}_1}{2}\right) \\ &< \frac{F(\mathbf{x}_0, \mathbf{y}_0) + F(\mathbf{x}_1, \mathbf{y}_1)}{2} \\ &= \frac{f(\mathbf{x}_0) + f(\mathbf{x}_1)}{2} \end{aligned}$$

which implies that f is strictly convex. 5.5

Theorem 5.6 Let $\bar{\ell} : \mathbb{R}^{|\lambda|} \longrightarrow (0, \infty]$ be defined by

$$\bar{\ell}(\mathbf{x}) = \min_{\mathbf{y} \in P_{\mathbf{x}}} L(\mathbf{x}, \mathbf{y}).$$

Then $\bar{\ell}$ is strictly convex and

$$\ell = \bar{\ell} \circ s_{\lambda}.$$

Proof. To see that the minimum defining $\bar{\ell}$ exists we use Lemma 5.3. At the same time we will see that ℓ is the composition of s_{λ} and $\bar{\ell}$. The set $\mathcal{P}(B, f)$ contains the geodesic representative in the homotopy class of $f(\alpha)$ so

$$\begin{aligned} \ell(B, f) &= \min_{\beta \in \mathcal{P}(B, f)} \text{length}(\beta) \\ &= \min_{\beta \in \mathcal{P}(B, f)} L(s_{\lambda}(B, f), p_{\lambda}(\beta)) \\ &= \min_{\mathbf{y} \in P_{\mathbf{x}}} L(\mathbf{x}, \mathbf{y}) \\ &= \bar{\ell} \circ s_{\lambda}(B, f) \end{aligned}$$

where second two inequalities follow from Lemma 5.3 and $\mathbf{x} = s_{\lambda}(B, f)$.

Let W be a wedge in $\mathcal{W}$. Since the function d is strictly convex the function $(\mathbf{x}, \mathbf{y}) \mapsto d_W(\gamma_W^-(\mathbf{y}), \gamma_W^+(\mathbf{y}) + \gamma_W^+(\mathbf{x}))$ is convex on $\mathbb{R}^{|\lambda|} \times \mathbb{R}^{|\lambda_0|}$ and is strictly convex along any line through distinct points $(\mathbf{x}_0, \mathbf{y}_0)$ and $(\mathbf{x}_1, \mathbf{y}_1)$ were either $\gamma_W^+(\mathbf{x}_0) \neq \gamma_W^+(\mathbf{x}_1)$, $\gamma_W^+(\mathbf{y}_0) \neq \gamma_W^+(\mathbf{y}_1)$ or $\gamma_W^-(\mathbf{y}_0) \neq \gamma_W^-(\mathbf{y}_1)$. Therefore if L is not strictly convex along this line we must have $\beta(\mathbf{x}_0) = \beta(\mathbf{x}_1)$ and $\beta(\mathbf{y}_0) = \beta(\mathbf{y}_1)$ for all $\beta \in \lambda_0$. For all closed curves γ in λ we then have $a_{\gamma}^{\pm}(\mathbf{x}_0, \mathbf{y}_0) = a_{\gamma}^{\pm}(\mathbf{x}_1, \mathbf{y}_1)$ and therefore if $\mathbf{y}_0 \in P_{\mathbf{x}_0}$ and $\mathbf{y}_1 \in P_{\mathbf{x}_1}$ we also have $\gamma(\mathbf{x}_0) = \gamma(\mathbf{x}_1)$. Therefore if L is not strictly convex along the line it can't lie in $\mathbb{P}$; equivalently L is strictly convex on $\mathbb{P}$. Lemma 5.5 then implies that $\bar{\ell}$ is strictly convex. 5.6

6 Proof of Proposition 3.6

We now return to a finite area hyperbolic surface X with an ideal triangulation λ and recall the proposition.

Proposition 3.6 *Let α be a closed curve on X and $\ell_\alpha : \mathcal{T}(X) \rightarrow \mathbb{R}^+$ its length function. Then there is a convex function $\bar{\ell}_\alpha : \mathbb{R}^{|\lambda|} \rightarrow \mathbb{R}$ such that $\ell_\alpha = \bar{\ell}_\alpha \circ s_\lambda$. Furthermore the $\bar{\ell}_\alpha$ -image of T_λ is in $\mathbb{R}^+$ and if α intersects every leaf of λ then $\bar{\ell}_\alpha$ is strictly convex.*

Let $\pi_\alpha : A_\alpha \rightarrow X$ be the annular covering such that α lifts to A_α . If (Y, f) is a pair in $\mathcal{T}(X)$ then $Y_{f(\alpha)}$ is the annulus cover of Y such that $f(\alpha)$ lifts to $Y_{f(\alpha)}$. The map $f : X \rightarrow Y$ lifts to a quasi-conformal homeomorphism $f_\alpha : X_\alpha \rightarrow Y_{f(\alpha)}$. This induces a map $\Pi_\alpha : \mathcal{T}(X) \rightarrow \mathcal{T}(X_\alpha)$. Let $\ell_\alpha : \mathcal{T}(X) \rightarrow (0, \infty)$ be the geodesic length function for α and let $\ell : \mathcal{T}(X_\alpha) \rightarrow (0, \infty)$ be the geodesic length function for the core curve of the annulus. Then $\ell_\alpha = \ell \circ \Pi_\alpha$.

Let λ_α be the set of geodesics in $\pi_\alpha^{-1}(\lambda)$ that intersect the core curve α . Then λ_α is an ideal wedgelation of X_α by Lemma 5.2. The covering map π_α defines a map $\bar{\pi}_\alpha$ from the leaves of λ_α to the leaves of λ which induces a map $\bar{\Pi}_\alpha : \mathbb{R}^{|\lambda|} \rightarrow \mathbb{R}^{|\lambda_\alpha|}$.

Proposition 6.1 *We can choose the coordinates for $\mathcal{T}(A)$ and $\mathcal{T}(S)$ such that $s_{\lambda_\alpha} \circ \Pi_\alpha = \bar{\Pi}_\alpha \circ s_\lambda$.*

Proof. The covering map $\pi : \mathbb{H}^2 \rightarrow X$ factors through π_α . In particular X and A_α have the same universal cover and the pre-image $\tilde{\lambda}_\alpha$ of λ_α is a subset of the pre-image $\tilde{\lambda}$ of λ . Assume we have chosen the coordinate map s_λ . Let γ' be a geodesic in λ_α and let $\gamma = \pi_\alpha(\gamma')$ be its image in λ . Let $\tilde{\gamma}'$ be a component of the pre-image of γ' in $\mathbb{H}^2$. In our choice of coordinate map s_λ we have chosen a pre-image $\tilde{\gamma}$ in $\mathbb{H}^2$ of γ and ideal triangles Δ_γ^a and Δ_γ^b . Then there is a deck transformation $g \in \pi_1(X)$ such that $g(\tilde{\gamma}) = \tilde{\gamma}'$. Since $\tilde{\gamma}$ separates Δ_γ^a from Δ_γ^b we have that $\tilde{\gamma}'$ separates $g(\Delta_\gamma^a)$ from $g(\Delta_\gamma^b)$. These ideal triangles will also be asymptotic to $\tilde{\gamma}'$ so they must have exactly two edges that intersect $\tilde{\gamma}'$, the pre-image of α in $\mathbb{H}^2$ and these edges will be in $\tilde{\lambda}_\alpha$. Exactly one of these ideal triangles, say $g(\Delta_\gamma^a)$, will be on the left $\tilde{\gamma}'$. Since it has two edges in $\tilde{\lambda}_\alpha$ there will be a wedge W in $\mathbb{H}^2 \setminus \tilde{\lambda}_\alpha$ with $\Delta(W) = g(\Delta_\gamma^a)$. We let $W_\gamma^- = W$. Similarly the ideal triangle $g(\Delta_\gamma^b)$ determines a wedge in $\mathbb{H}^2 \setminus \tilde{\lambda}_\alpha$ which we define to be W_γ^+ . If $g(\Delta_\gamma^b)$ is on the left we reverse the labels. It is then straightforward to check that the lemma holds. 6.1

Let $\bar{\ell} : \mathbb{R}^{|\lambda_\alpha|} \rightarrow [0, \infty]$ be the factorization of ℓ given in the previous subsection so that $\ell = \bar{\ell} \circ s_{\lambda_\alpha}$. Define $\bar{\ell}_\alpha : \mathbb{R}^{|\lambda|} \rightarrow [0, \infty]$ by $\bar{\ell}_\alpha = \bar{\ell} \circ \bar{\Pi}_\alpha$. We then have the following commutative diagram:

$$\begin{array}{ccccc}
T(X) & \xrightarrow{\sigma_\lambda} & \mathbb{R}^{|\lambda|} & & \\
\downarrow \Pi_\alpha & \searrow \ell_\alpha & \swarrow \bar{\ell}_\alpha & & \downarrow \bar{\Pi}_\alpha \\
& & [0, \infty] & & \\
\uparrow \ell & \nwarrow \bar{\ell} & \swarrow \bar{\ell}_\alpha & & \uparrow \bar{\Pi}_\alpha \\
T(X_\alpha) & \xrightarrow{\sigma_{\lambda_\alpha}} & \mathbb{R}^{|\lambda_\alpha|} & &
\end{array}$$

The first claim of Proposition 3.6 follows directly from the commutativity of the triangle on the top. The convexity of $\bar{\ell}_\alpha$ follows from the strict convexity of $\bar{\ell}$ and the fact that $\bar{\Pi}_\alpha$ is linear. In the case that α intersects every leaf of λ , the map $\bar{\Pi}_\alpha$ is also injective and hence $\bar{\ell}_\alpha$ is convex as claimed. The following lemma now concludes the proof of Proposition 3.6:

Lemma 6.2 *The function $\bar{\ell}_\alpha$ is finite on T_λ .*

If we knew at this stage that s_λ is a homeomorphism onto its image T_λ , the claim of Lemma 6.2 would follow also from the commutativity of the diagram above and the fact that ℓ_α is obviously finite. Unfortunately, we derived this fact from Proposition 3.6.

Proof. Let $\mathbf{x}$ be in T_λ and let $\mathbf{x}_\alpha = \bar{\Pi}_\alpha(\mathbf{x})$. We will choose a $\mathbf{y} \in \mathbb{R}^{|(\lambda_\alpha)_0|}$ such that $\mathbf{y} \in \mathcal{P}_{\mathbf{x}_\alpha}$ and $L(\mathbf{x}_\alpha, \mathbf{y}) < \infty$. Let γ_α be a non-isolated leaf of λ_α . Then $\bar{\pi}_\alpha(\gamma_\alpha)$ is a closed leaf γ of λ . Let γ_i^+ be the leaves on the right of γ_α labeled as the were in the previous section. Assume that the right side of γ_α maps to the a -side of γ . Then $\bar{\pi}_\alpha(\gamma_i^+) = \bar{\pi}_\alpha(\gamma_{i+k}^+)$ where

$$k = \sum_{\beta \in \lambda} n_\gamma^a(\beta).$$

Therefore $\gamma_i^+(\mathbf{x}_\alpha) = \gamma_{i+k}^+(\mathbf{x}_\alpha)$ and there is a constant c such that

$$c = \sum_{i=j}^{j+k-1} \gamma_i^+(\mathbf{x}_\alpha)$$

for all $j \geq 0$. It follows that

$$\sum_{i=0}^{nk+j} \gamma_i^+(\mathbf{x}_\alpha) = nc + \sum_{i=0}^j \gamma_i^+(\mathbf{x}_\alpha).$$

Choose the γ_i^+ -coordinates of $\mathbf{y}$ such that

$$\gamma_n^+(\mathbf{y}) = \sum_{i=0}^{n-1} \gamma_i^+(\mathbf{x}_\alpha) = \gamma_{n-1}^+(\mathbf{y}) + \gamma_{n-1}^+(\mathbf{x}_\alpha).$$

Let $\mathcal{W}_{\gamma_\alpha}^+$ be the set of wedges in $\mathcal{W}$ whose boundary edges are in γ_i^+ and γ_i^+ for some $i \geq 0$. Then

$$\begin{aligned}
\sum_{W \in \mathcal{W}_{\gamma_\alpha}^+} d_W(\gamma_W^-(\mathbf{y}), \gamma_W^+(\mathbf{y}) + \gamma_W^+(\mathbf{x}_\alpha)) &= \sum_{j=0}^{k-1} \sum_{i=0}^{\infty} d\left(\gamma_{j+ki+1}^+(\mathbf{y}), \gamma_{j+ki}^+(\mathbf{y}) + \gamma_{j+ki}^+(\mathbf{x}_\alpha)\right) \\
&= \sum_{j=0}^{k-1} \sum_{i=0}^{\infty} d\left(ic + \gamma_j^+(\mathbf{x}_\alpha), ic + \gamma_j^+(\mathbf{x}_\alpha)\right) \\
&\leq \sum_{j=0}^{k-1} \sum_{i=0}^{\infty} e^{-ic - \gamma_j^+(\mathbf{x}_\alpha)} \\
&< \infty
\end{aligned}$$

where the inequality on the third line follows from Lemma 5.1.

We now take the leaves γ_i^- on the left of γ_α . We similarly choose the γ_i^- -coordinates but so that we have $a_{\gamma_\alpha}^+(\mathbf{x}_\alpha, \mathbf{y}) + \gamma_\alpha(\mathbf{x}_\alpha) = a_{\gamma_\alpha}^-(\mathbf{x}_\alpha, \mathbf{y})$ we define

$$\gamma_n^-(\mathbf{y}) = \gamma_\alpha(\mathbf{x}_\alpha) - \sum_{i=0}^n \gamma_i^-(\mathbf{x}_\alpha).$$

Similar to above we define $\mathcal{W}_{\gamma_\alpha}^-$ to be the set of wedges in $\mathcal{W}$ whose boundary edges are γ_i^- and γ_{i+1}^- for some $i \geq 0$ we similarly find that

$$\sum_{W \in \mathcal{W}_{\gamma_\alpha}^-} d_W(\gamma_W^-(\mathbf{y}), \gamma_W^+(\mathbf{y}) + \gamma_W^+(\mathbf{x}_\alpha)) < \infty.$$

We repeat the above construction for each of the finitely many non-isolated leaves of λ_α . Note that each wedge $W \in \mathcal{W}$ is asymptotic to at most one of the non-isolated leaves γ_α and therefore is in at most one of the sets $\mathcal{W}_{\gamma_\alpha}^+$ or $\mathcal{W}_{\gamma_\alpha}^-$ that contains W and all but finitely many of the W are in one of these sets. This defines $\mathbf{y}$ for all but finitely many coordinates. The remaining we choose freely. We can then split the infinite sum $L(\mathbf{x}_\alpha, \mathbf{y})$ into finitely many sums each of which is finite. Therefore $\bar{\ell}_\alpha(\mathbf{x}) \leq L(\mathbf{x}_\alpha, \mathbf{y}) < \infty$. 6.2

This concludes the proof of Proposition 3.6 and with it the whole paper.

References

- [1] F. Bonahon, *Shearing hyperbolic surfaces, bending pleated surfaces and Thurston's symplectic form*, Ann. Fac. Sci. Toulouse Math. (6) 5 (1996).

- [2] P. Buser, *Geometry and spectra of compact Riemann surfaces*, Progress in Mathematics, 106. Birkhäuser, 1992.
- [3] A. Casson and S. Bleiler, *Automorphisms of surfaces after Nielsen and Thurston*, London Mathematical Society Student Texts, 9. Cambridge University Press, 1988.
- [4] A. Casson, and D. Jungreis, *Convergence groups and Seifert fibered 3-manifolds*, Invent. Math. 118 (1994).
- [5] D. Gabai, *Convergence groups are Fuchsian groups*, Ann. of Math. (2) 136 (1992).
- [6] Y. Iwayoshi and M. Taniguchi, *An introduction to Teichmüller spaces*, Springer-Verlag, Tokyo, 1992.
- [7] S. Kerckhoff, *The Nielsen realization problem*, Ann. of Math. (2) 117 (1983).
- [8] S. Wolpert, *Geodesic length functions and the Nielsen problem*, J. Differential Geom. 25 (1987)
- [9] W. Thurston, *Hyperbolic Structures on 3-manifolds, II: Surface groups and 3-manifolds which fiber over the circle*, arXiv [math.GT/9801045](https://arxiv.org/abs/math.GT/9801045)
- [10] A. Tromba, *Dirichlet's energy on Teichmüller's moduli space and the Nielsen realization problem*, Math. Z. 222 (1996).

K. Bromberg
 Department of Mathematics, University of Utah
bromberg@math.utah.edu

M. Bestvina
 Department of Mathematics, University of Utah
bestvina@math.utah.edu

K. Fujiwara
 Graduate School of Information Sciences, Tohoku University
fujiwara@math.is.tohoku.ac.jp

J. Souto
 Department of Mathematics, University of Michigan
jsouto@umich.edu